

## DOCUMENT RESUME

ED 460 531

EC 306 907

AUTHOR Costello, Elizabeth Jane; Horwitz, Sarah McCue; Angold, Adrian; Hoagwood, Kimberly; Sparrow, Sara S.; Triche, Elizabeth

TITLE Symptoms and Functioning: Service Implications of Recent Research on Serious Emotional Disturbance. Symposium.

PUB DATE 1996-02-00

NOTE 6p.; In: A System of Care for Children's Mental Health: Expanding the Research Base. Proceedings of the Annual Research Conference (9th, Tampa, FL, February 26-28, 1996); see EC 306 844.

AVAILABLE FROM For full text:  
<http://rtckids.fmhi.usf.edu/Proceed9th/9thprocindex.htm>.

PUB TYPE Collected Works - General (020) -- Reports - Research (143) -- Speeches/Meeting Papers (150)


EDRS PRICE MF01/PC01 Plus Postage.

DESCRIPTORS \*Accessibility (for Disabled); \*Adaptive Behavior (of Disabled); Compliance (Legal); Eligibility; \*Emotional Disturbances; \*Federal Regulation; Incidence; Mental Health Programs; Research and Development; Rural Areas; \*Theory Practice Relationship; Validity

## ABSTRACT

Three brief papers review the literature and report on a study concerning the concept of functional impairment in light of regulations requiring both a psychiatric diagnosis and functional impairment for access to mental health care by children and adolescents with emotional disturbances. The first, very brief summary is: "The Origin, Meaning and Use of Adaptive Functioning in Defining Mental Health Problems" (Sara M. Horwitz and others), which traces the history of the concept of adaptive functioning. The second summary is: "Service Needs and Use for Serious Emotional Disturbance: A Community Study" (E. Jane Costello). This study examined the prevalence of serious emotional disturbance (SED) in a large rural area and found that SED criteria identify about one child in 10, of which only one in three is currently receiving any services. The third summary is: "The Validity of Serious Emotional Disturbance as a Marker for Need for Services" (Adrian Angold). It finds that while the requirement of both psychiatric diagnosis and functional impairment does identify children with high levels of need for service, many children with marked functional impairment but less severe psychopathology will be excluded from services. (DB)

ED 460 531

	<p style="text-align: center;"><b>Symposium:</b></p> <p style="text-align: center;"><b>Symptoms and Functioning: Service Implications of Recent Research on Serious Emotional Disturbance</b></p> <hr/> <p style="text-align: center;"><u>Authors</u></p> <p style="text-align: center;"><u>Introduction</u> <u>Origin, Meaning and Use</u> <u>Service Needs</u> <u>Validity</u></p>
<p><b>Shortcuts</b></p>	<p><b>Introduction</b></p> <p>E. Jane Costello</p> <p>Recent changes in regulations governing access to mental health care resources for children have included requirements that children not only have a psychiatric disorder but also be functionally impaired. This has aroused interest in the concept of functional impairment or adaptive functioning (i.e., its history, measurement, and significance as an indicator of need for services). In this symposium, (a) Dr. Horwitz traces the history of the concept, (b) Dr. Costello describes its prevalence and correlates in the community, and (c) Dr. Angold discusses evidence for impairment as a marker of service need.</p>

U.S. DEPARTMENT OF EDUCATION  
Office of Educational Research and Improvement  
EDUCATIONAL RESOURCES INFORMATION  
CENTER (ERIC)

☒ This document has been reproduced as  
received from the person or organization  
originating it  
☐ Minor changes have been made to improve  
reproduction quality

• Points of view or opinions stated in this docu-  
ment do not necessarily represent official  
OERI position or policy

EC 306 907

## **The Origin, Meaning and Use of Adaptive Functioning in Defining Mental Health Problems**

Sara M. Horwitz, K. Hoagwood, S. S. Sparrow & E. Triche

The concept of adaptive functioning grew out of the study of mental retardation. Concerns with decreased intellectual functioning became important because of increased societal demands brought on by industrialization, urbanization, and universal public education. Early work around quantification of intellectual abilities revealed variations in functioning not uniformly related to intellectual levels. These differences in functioning were related, however, to an individual's ability to perform age-appropriate social roles and became incorporated into the definition of mental retardation. For those with severe and persistent mental illness, the need to differentiate individuals with respect to prognosis and services planning led to an interest in adaptive functioning and its eventual inclusion in the multiaxial classification system of the Diagnostic and Statistical Manual (DSM), 3rd Edition. Unfortunately, measurement of adaptive functioning within the definition of mental disorders is underdeveloped and has been confined largely to a single global measure of functional impairment.

## **Service Needs and Use for Serious Emotional Disturbance: A Community Study**

E. Jane Costello

Many federal and state agencies and insurance companies are adopting the federal definition of Serious Emotional Disturbance (SED) as the criterion for child psychopathology of sufficient severity to require services. SED is defined as a DSM-defined psychiatric disorder resulting in functional impairment that substantially interferes with or limits the child's role or functioning in family, school, or community activities.

### **Methods**

In the Great Smoky Mountains Study, we examined the prevalence of SED in a representative population sample of children age 9 - 13, randomly selected from a largely rural area of the southeastern United States. Interviews with 1,015 parent and child pairs were conducted by trained interviewers using (a) the Child and Adolescent Psychiatric Assessment (CAPA) which assesses DSM-III-R disorders and functional impairment, and (b) the Child and Adolescent Services Assessment (CASA) which records service use across five service sectors (i.e., specialty mental health, education, primary care, juvenile justice, and child welfare). Functional impairment was also evaluated using the Children's Global Assessment Scale (C-GAS) and the North Carolina Child and Adolescent Functional

North Carolina Child and Adolescent Functional Assessment Scale (CAFAS). Prevalence estimates were adjusted for design effects.

## **Results**

Of the 20.3% of the population who had one or more DSM-III-R disorders, 55% (11.1% of the population) met criteria for SED according to one or more of the three measures of impairment used (i.e., CAPA, C-GAS, or CAFAS). Boys and girls were equally likely to have SED. Of children in the sample with conduct disorder, 78.8% had SED, compared with 73.4% of those with depressive disorders, 91.3% of those with oppositional defiant disorders, and 73.1% of those with ADHD. Disorders least likely to be associated with functional impairment were anxiety disorders (45.8%), tics, and functional enuresis (32.6% each).

Children with SED used services at a higher rate than those with only a DSM diagnosis or only functional impairment. However, the majority of children with SED had received no mental health services from any service sector during the three months preceding the interview. Of children with SED, 19.2% had received services in the specialty mental health sector, 29.1% in the school system, 6% from primary care providers, 5.3% through child welfare agencies, and 2% through juvenile justice. In total, only 37.7% of children with SED received mental health care from any agency, compared with 6% of children with only a DSM diagnosis or functional impairment.

## **Conclusions**

Use of the SED criterion will identify around one child in ten in the population, depending on the criteria used. These youth are much more likely than children with less severe symptoms to receive mental health care; however, only about one in three is currently receiving any form of care. Thus, adopting SED as the definition of need for care identifies a large gap between need and availability of services.

## **The Validity of Serious Emotional Disturbance as a Marker for Need for Services**

Adrian Angold

The addition of functional impairment to diagnosis as a criterion for eligibility for various treatment and reimbursement programs reflects an attempt to focus scarce resources on those in greatest need. In this paper we explore some empirical evidence testing the hypothesis that children with SED have greater service needs than children with (a) functional impairment associated with psychiatric symptoms but no specific DSM diagnosis, (b) children with impaired functioning but few or no psychiatric symptoms,

(c) children with a DSM disorder but no functional impairment, or (d) children with neither psychopathology nor impairment.

### **Method**

The Great Smoky Mountains Study's representative population sample of youth aged 9 - 13 (N = 1,015) was used for these analyses. Psychiatric symptoms were assessed by trained lay interviewers using the Child and Adolescent Psychiatric Assessment (CAPA), and service use across five service sectors was recorded using the Child and Adolescent Services Assessment (CASA). Service need was defined in the following ways: (a) use of specialty mental health services; (b) use of school counselors and school psychologists for emotional and behavioral problems; (c) parental report of economic, social, or psychological burdens associated with the child's symptoms; (d) report by parent or child that the child had a problem in one or more symptom areas; and (e) report by parent or child that help was needed because of such a problem.

### **Results**

Children with SED had the highest rates of use of specialty mental health and school services, the highest rates of burden, reports of problems, and need for help. However, children with functional impairment, with or without some psychiatric symptoms, were more likely than children with a DSM-III-R diagnosis but no functional impairment to meet the various criteria of need for services. Between the first and second years of the study, there was a considerable amount of movement between categories for severity of disorder.

### **Conclusions**

While the combination of a DSM diagnosis and functional impairment (SED) identifies children with high levels of need for services, service needs are also high in youth with less severe psychopathology but marked functional impairment. If these children are excluded from access to care because agencies adopt SED as the criterion for access to care, many children in need will be excluded. Furthermore, it will exclude from services children with less severe symptoms but high levels of functional impairment who are likely to move into full-blown SED without treatment.

## Authors

Elizabeth Jane Costello, Ph.D.  
Associate Professor  
Developmental Epidemiology Program  
Duke University Medical Center  
Box 3454  
Durham, NC 27710  
Voice: 919/687-4686  
Fax: 919/687-4737  
jcostell@psych.mc.duke.edu

Sarah McCue Horwitz, Ph.D.  
Associate Professor  
Department of Epidemiology and Public Health  
Yale University  
60 College Street  
New Haven, CT 06520  
Voice: 203/785-2862  
Fax: 203/785-6287  
kriegerpb@maspo3.mas.yale.edu

Adrian Angold, MRCPsych  
Assistant Professor  
Developmental Epidemiology Program  
Duke University Medical Center  
Box 3454  
Durham, NC 27710  
Voice: 919/687-4686  
Fax: 919/687-4737  
ztbrooks@psych.mc.duke.edu

Kimberly Hoagwood, Ph.D.  
Child/Adolescent Mental Health Services Branch  
National Institute of Mental Health  
5600 Fishers Lane, Parklawn Building, Room 10C-06  
Rockville, MD 20857  
Voice: 301/443-4233

Sara S. Sparrow, Ph.D.  
Yale University School of Medicine  
Child Study Center, 1172 SHM  
230 South Frontage Road  
New Haven, CT 06520  
Voice: 203/785-6227

Elizabeth Triche, M.Phil.  
Department of Epidemiology & Public Health  
60 College Street  
New Haven, CT 06520  
Voice: 203/785-2862

Liberton, C., Kutash, K., & Friedman, R. M. (Eds.), (1997). The 9th Annual Research Conference Proceedings, A System of Care for Children's Mental Health: Expanding the Research Base (February 26 - 28, 1996). Tampa, FL: University of South Florida, Louis de la Parte Florida Mental Health Institute, Research & Training Center for Children's Mental Health.



**U.S. Department of Education**  
Office of Educational Research and Improvement (OERI)  
National Library of Education (NLE)  
Educational Resources Information Center (ERIC)



## **NOTICE**

### **REPRODUCTION BASIS**



This document is covered by a signed "Reproduction Release (Blanket) form (on file within the ERIC system), encompassing all or classes of documents from its source organization and, therefore, does not require a "Specific Document" Release form.



This document is Federally-funded, or carries its own permission to reproduce, or is otherwise in the public domain and, therefore, may be reproduced by ERIC without a signed Reproduction Release form (either "Specific Document" or "Blanket").